

Reply to the Comment “Intercluster correlation in Seismicity” by Helmstetter and Sornette

The comment by Helmstetter and Sornette [1] addresses an important point of our paper [2] that needs to be clarified. Any scaling measure, including also the diffusion entropy (DE) method, when applied to the earthquake time series, yields anomalous scaling for two reasons, long-range time correlation (intermittency) and the Pareto’s statistics. In fact, the DE random walker position is determined by the number of events (earthquakes) in a given time interval, which, in turns, is related to both the number of clusters and the cluster size. The distribution of the time distances between clusters, $\psi(\tau^{[m]})$ in ref. [2], determines the number of clusters, while the Pareto’s law

$$p(n) \propto n^{-\gamma} \quad (1)$$

gives the probability to get a cluster of size n . The resulting asymptotic scaling detected by the DE is the scaling of the most anomalous of the two processes, i.e. the process with the lowest power law exponent.

The authors of the comment [1] discuss the Generalized Poisson (GP) model with an exponent $\gamma = 2.25$ [3], which generates $\delta = 0.8$. In fact, according to the Continuous Time Random Walk theory [4], the scaling is $\delta = 1/(\gamma - 1)$ if $2 < \gamma < 3$, $\delta = 0.5$ for $\gamma > 3$ and $\delta = 1$ for $\gamma < 2$.




FIG. 1. We plot the DE analysis for different synthetic series generated with the GP model and three different exponents of the Pareto law, i.e. $\gamma = 3.5, 2.25, 1.25$. The values of δ obtained by fitting the asymptotic behavior are also reported with the corresponding error. These values increase as γ diminishes passing respectively from 0.52 ± 0.02 (for $\gamma = 3.5$) to 0.78 ± 0.02 (for $\gamma = 2.25$) and finally to 0.99 ± 0.02 (for $\gamma = 1.25$).

We have repeated the calculations with $\gamma = 2.25$ and the results, shown in fig.1, confirm the theoretical prediction. In the figure we plot also two other cases in order to illustrate how δ changes as a function of γ . In Ref. [2] we explored the range $\gamma = 2.5 - 3$.

The experimental data yield a value $\delta = 0.94$ confirming that $\psi(\tau^{[m]})$ has a power law tail with an exponent $\mu = 2.06$. Moreover, the fact that the scaling observed in fig. 2 of our previous paper [2] is not influenced by the threshold of magnitude considered, is an additional evidence that what we are actually observing is the real scaling determined by the clusters distances.

The LR model [2] is a minimal generalization of the GP model (only $\psi(\tau^{[m]})$ changes). Therefore it is very interesting that a more sophisticated model, like the ETAS, which reproduces also other properties of seismicity, yields the same δ observed in the LR and in the experimental data. However we would like to point out that we define $\tau^{[m]}$ as the distance between two consecutive clusters, while in [1] this symbol refers to distance between two consecutive earthquakes with magnitude greater than $M = 5$. So there is no contradiction between the correlations found in [1] and our results.

Mirko S. Mega¹, Paolo Allegrini², Paolo Grigolini^{1,3,4}, Vito Latora⁵, Luigi Palatella¹, Andrea Rapisarda⁵ and Sergio Vinciguerra^{5,6}

¹Dipartimento di Fisica dell’Università di Pisa and INFN, via Buonarroti 2, 56127 Pisa, Italy

²Istituto di Linguistica Computazionale del CNR, Area della Ricerca di Pisa, Via G. Moruzzi 1, 56124, Pisa, Italy

³Center for Nonlinear Science, University of North Texas, P.O. Box 311427, Denton, Texas 76203-1427

⁴Istituto di Biofisica del CNR, Area della Ricerca di Pisa, Via Alfieri 1, San Cataldo, 56010, Ghezzano-Pisa, Italy

⁵Dipartimento di Fisica e Astronomia, Università di Catania, and INFN sezione di Catania, Via S. Sofia 64, 95123 Catania, Italy

⁶Osservatorio Vesuviano - INGV, Via Diocleziano 328, 80124 Napoli, Italy

PACS numbers: 91.30.Dk, 05.45.Tp, 05.40.Fb

[1] comment by A. Helmstetter and D. Sornette, physics/0307134

[2] M. S. Mega, P. Allegrini, P. Grigolini, V. Latora, L. Palatella, A. Rapisarda and S. Vinciguerra, Phys. Rev. Lett. **90**, 188501 (2003).

[3] In the literature Pareto’s law is referred as the cumulative distribution. Our exponent γ is related to the exponent reported in ref. [1] by the expression $\gamma = 1 + b/\alpha$.

[4] P. Grigolini, L. Palatella, G. Raffaelli, Fractals **9**, 439 (2001).